

# Improving Refill Adherence in Medicare Patients With Tailored and Interactive Mobile Text Messaging

AUTHORED BY



**KAISER  
PERMANENTE®**



PUBLISHED BY

**JMIR mHealth  
and uHealth**



Original Paper

# Improving Refill Adherence in Medicare Patients With Tailored and Interactive Mobile Text Messaging: Pilot Study

Rena Brar Prayaga<sup>1</sup>, MA, JD; Erwin W Jeong<sup>2</sup>, PharmD; Erin Feger<sup>1</sup>, BA; Harmony K Noble<sup>3</sup>, BA; Magdalen Kmiec<sup>1</sup>, BA; Ram S Prayaga<sup>1</sup>, MS

<sup>1</sup>mPulse Mobile, Inc, Encino, CA, United States

<sup>2</sup>Medicare Medication Therapy Management, Kaiser Permanente Southern California, Downey, CA, United States

<sup>3</sup>Regional Pharmacy Clinical Operations, Kaiser Permanente Southern California, Downey, CA, United States

**Corresponding Author:**

Rena Brar Prayaga, MA, JD

mPulse Mobile, Inc

16530 Ventura Blvd. Suite 500

Encino, CA,

United States

Phone: 1 888 678 5735

Email: [rena@mpulsemobile.com](mailto:rena@mpulsemobile.com)

## Abstract

**Background:** Nonadherence is a major concern in the management of chronic conditions such as hypertension, cardiovascular disease, and diabetes where patients may discontinue or interrupt their medication for a variety of reasons. Text message reminders have been used to improve adherence. However, few programs or studies have explored the benefits of text messaging with older populations and at scale. In this paper, we present a program design using tailored and interactive text messaging to improve refill rates of partially adherent or nonadherent Medicare members of a large integrated health plan.

**Objective:** The aim of this 3-month program was to gain an understanding of whether tailored interactive text message dialogues could be used to improve medication refills in Medicare patients with one or more chronic diseases.

**Methods:** We used the mPulse Mobile interactive text messaging solution with partially adherent and nonadherent Medicare patients (ie, over age 65 years or younger with disabilities) of Kaiser Permanente Southern California (KP), a large integrated health plan, and compared refill rates of the text messaging group (n=12,272) to a group of partially adherent or nonadherent Medicare patients at KP who did not receive text messages (nontext messaging group, n=76,068). Both groups were exposed to other forms of refill and adherence outreach including phone calls, secure emails, and robo-calls from December 2016 to February 2017.

**Results:** The text messaging group and nontext messaging group were compared using an independent samples t test to test difference in group average of refill rates. There was a significant difference in medication refill rates between the 2 groups, with a 14.07 percentage points higher refill rate in the text messaging group ( $P<.001$ ).

**Conclusions:** The results showed a strong benefit of using this text messaging solution to improve medication refill rates among Medicare patients. These findings also support using interactive text messaging as a cost-effective, convenient, and user-friendly solution for patient engagement. Program outcomes and insights can be used to enhance the design of future text-based solutions to improve health outcomes and promote adherence and long-term behavior change.

(*JMIR Mhealth Uhealth* 2018;6(1):e30) doi:[10.2196/mhealth.8930](https://doi.org/10.2196/mhealth.8930)

**KEYWORDS**

patient activation; patient engagement; medication adherence; refill management; text messaging; interactive; NLP; Medicare; disease management; technology acceptability model

## Introduction

### Overview

Patient nonadherence affects 50% to 60% of chronically ill patients, and the cost of medication-related hospitalizations is \$100 billion annually [1-3]. It is also associated with poor outcomes and progression of disease causing approximately 125,000 deaths and at least 10% of hospitalizations every year [4]. Seniors take an average of 7 medications per day, representing the highest number of prescribed medications for any age group [5].

Nonadherence is a major concern in the management of chronic conditions such as hypertension, cardiovascular disease, and diabetes where patients may discontinue or interrupt their medication for a variety of reasons. Patients are considered adherent when they take their medications (dose, time, frequency) as prescribed by their health care provider and as agreed to by the patient. Medicare populations adherence rates are often measured by pharmacy refill rates. The Centers for Medicare and Medicaid Services (CMS) uses the proportion of days covered (PDC), developed by Pharmacy Quality Alliance, to calculate adherence. Based on this, a patient who has a PDC rate of at least 80% is considered to be adherent.

Adherence is a particularly difficult problem among Medicare populations, and adherence rate is a key metric used by CMS to measure quality of a managed care plan. Approximately 32% of Medicare Part D patients are nonadherent to their diabetes, hypertension, and cholesterol medications [6]. Reasons for nonadherence may include side effects of the drug, cost of the drug, lack of perceived benefit, and/or forgetfulness.

### Use of Mobile Technology for Adherence

Studies and surveys are finding that digital health is not reaching most seniors, especially where there are socioeconomic disparities [7]. Among seniors who are identified as tech-savvy, 70% of those surveyed believe it's important to be able to request prescription refills electronically, but fewer than half (46%) say they can do so today [8]. On researching mobile phone device ownership among seniors, we learned that while 78% of Americans aged 65 years and older own a mobile phone, only 34% own a smartphone [9,10]. We estimated smartphone ownership to be even lower among Medicare populations aged 65 years and older.

Text messaging using SMS (short message service) is ubiquitous, highly accessible, affordable, and commonly used across all income levels. It is also an effective channel for continuing to engage individuals in their health care once they leave the doctor's office. Interactive text dialogues provide an opportunity for patients and health plan members to tap into health care resources and get support for healthy behaviors and long-term behavior change. Several studies have found that text messaging may serve as a scalable and effective means to improve medication adherence in chronic disease populations [11,12]. While there has been an interest in developing health technologies such as reminder applications [13-16] or automated phone reminders for older populations [17], a review of the literature reveals that very few programs have explored using

text messaging with seniors to improve medication refill adherence [18,19].

We determined at the outset that since the target users for the program were an older and/or disabled population on Medicare, it would be important to focus on usability (ie, ease of use) and simplicity (ie, design for basic feature mobile phone instead of smartphone). We used Davis' technology acceptance model (TAM) [20] as a guide to predict and optimize user acceptance of our solution as a viable and dynamic channel for interactive communication [21]. Therefore, our content strategy focused on usefulness and ease of use by providing simple instructions for authentication and task completion [22].

### Objectives

The program objectives were to assess the impact of an interactive and easy-to-use text messaging solution on medication refills and pharmacy operations and efficiencies. The target population consisted of partially adherent and nonadherent Medicare patients of a large integrated health plan (Kaiser Permanente Southern California, or KP) with 1 or more chronic diseases.

Our hypothesis was that patients receiving text message refill reminders (text messaging group) in addition to existing outreach would have a higher medication refill rate compared to the group that did not receive text messages (nontext messaging group).

## Methods

### Participants

The program began on December 7, 2016. All patients were Medicare members of KP with 1 or more chronic conditions (diabetes, hypertension, and/or high cholesterol). Patients in this program would be refilling 1 or more of the following 3 classes of drugs: oral diabetes medications, blood pressure medications (renin-angiotensin system antagonists), and statins.

There were approximately 5000 to 14,000 patients each week on the list who required pharmacy follow-up. These patients were pulled from 3 separate KP lists: (1) New Start: patients who filled their medication the first time in the calendar year and had a day supply remaining (DSR) of 0 to 30 days, (2) Near Goal: patients whose DSR ranged from -7 to 7 days and PDC ranged from 78% to 85%, and (3) Nonadherent: patients who had 2 fills within the calendar year and need to refill their medication by a specific date (Nonadherent date) in order to have a chance to improve their PDC to 80% or higher. The Nonadherent list patients were messaged in month 1 (December 2016) only.

Patients were divided into 2 groups:

1. Text message group (12,272/88,340, 13.89%): those who had opted in to receive text messages (as recorded within the health system's electronic medical records [EMR]) and were on the weekly list for pharmacy follow-up (1000 to 4000 patients per week). These patients received text messages reminding them to refill their prescriptions. This group consisted of 12,272 patients who had opted in to receive text messages and did in fact receive text messages

over the course of the program. [Table 1](#) provides age and race/ethnicity breakdowns for this group.

2. Nontext message group (76,068/88,340, 86.11%): those who had not opted in to receive text messages or there was no indication of an opt-in (as recorded within the health system's EMR) and were on the weekly list for pharmacy follow-up (4000 to 10,000 patients per week). This group consisted of 76,068 patients who did not receive text messages over the course of the program.

The text messaging group was one-fifth the size of the nontext messaging group because we were targeting only those Medicare patients who had opted in to receive text messages from KP. Both groups also received usual care which included phone calls and/or robo-calls reminding them to refill their prescriptions.

The Kaiser Permanente Southern California Institutional Review Board determined that this program did not involve human subject research and review was not necessary.

## Procedure

### Solution Overview

The mPulse Mobile platform delivers text messages to patients and members on behalf of health care companies. The platform


consists of several components that together enable companies to interactively engage with their end-users about appointments, refills, gaps in care, or other health-related topics. Patients in the text messaging group received a refill reminder dialogue that consisted of a series of messages. All messages were written at a 6th grade readability level. The first message was a greeting reminding them that they were due for a refill. They were then prompted to enter their date of birth to authenticate and view their refill order ([Figure 1](#)).


Upon confirmation of the order by the patient, the KP pharmacy received a notification, and a KP pharmacist would process the refill and use the mPulse Engagement Console to inform the patient when it would be ready for pickup. Patients who did not respond to the initial message in the dialogue would receive follow-up reminders 2 hours later and again 24 hours later. They would then be moved through the same process (authentication, confirming refill order, etc). After confirmation of the order, there was no further communication with the patient. However, a small subset of patients was messaged again in a later dialogue because they failed to refill again the following month. A more detailed view of the dialogues and the process is provided in [Multimedia Appendix 1](#).

**Table 1.** Characteristics of text messaging group.

Characteristic	Value, %
<b>Age, years</b>	
Under 65	13.2
65-70	39.7
70-75	24.1
75-85	18.9
Over 85	4.1
<b>Race/ethnicity</b>	
White	41.6
Hispanic/Latino	30.0
Black/African American	14.7
Asian/native Hawaiian	10.9
Unknown	2.75

**Figure 1.** Overview of message flow within refill dialogue.



**Figure 2.** Engagement Console used to process refill requests and address other concerns via text.


The screenshot displays the 'Mobile Engagement Console' interface. At the top, there's a header with the mPulse Mobile logo and navigation links: 'CONTACT MANAGER', 'CALL CENTER', and 'LOG OUT'. Below the header, a filter bar shows 'All Call Status', 'San Diego', 'All Actions', '2017-01-02 to 2017-02-01', and 'Export Members'. The main area features a table with columns: 'Call Status', 'Close Workflow', 'Service Area', 'Action Bucket', 'Name', 'Phone Number', and 'Date Added'. The first row is highlighted in green, showing 'Calling', 'Close Workflow', 'San Diego', 'Free Text', and a timestamp of '02/01/2017 12:56:47'. Below this, a detailed view of a contact is shown, including a 'View Contact Manager Profile' link, source file, drug name (Lisinopril-Hydrochlorothiazide 20-12.5 Mg Tabs), birth date (07/25/1936), drug class (Ras), and pharmacy (Central Mail Order Pharmacy). A list of actions follows, with columns for 'Open', 'Close Workflow', 'Service Area', 'Action Bucket', 'Name', 'Phone Number', and 'Date Added'. The actions include 'Open', 'Change', 'Incomplete', 'Free Text', and 'Dob' for various dates in February 2017.

Patients could move through the dialogue and authenticate their date of birth, complete a refill, ask for help, share reasons why they had not refilled already, or choose to opt out by using numeric or textual responses on their phone. The simplicity of the process allowed older users, who might also be more likely to have mobile phones instead of smartphones, to express their preferences and complete the process very easily.

If patients responded that they were experiencing side effects; did not believe the medication was helping them; wanted to change their medication, dose, or pharmacy; or had other concerns that might require follow-up, mPulse Mobile sent a daily list of members with pending questions or issues to the KP pharmacy for follow-up.

### Dialogue Initiation

Refill dialogues were initiated at 10 am every Wednesday and Thursday to allow for a reasonable time frame within which patients could respond. Patients who texted STOP or 7867 (easier option for those without smartphones) would be opted out from the campaign and would not receive any further messages. Dialogues included tailored information to customize the message content (eg, name, date of birth, drug, pharmacy).

Patient information such as phone number, drug names, gender, name, mobile opt-in, level of adherence, and date of birth was used in 2 ways: to tailor message content for patients and initiate reminder dialogues to patients based on exclusion and combination logic. This logic helped avoid duplication and over-messaging (eg, member on multiple lists or multiple drugs would still receive a single dialogue). Patient information was provided weekly from the integrated health system and was used to perform dialogue assignments every week.

### Refill Requests and Processing

Refill requests, questions, and concerns were handled by the pharmacy staff with a total of 8 staff members being trained on how to use the Engagement Console. To process refill requests or other concerns, staff would log on to a personalized view of the Engagement Console (based on their assigned medical center) and would be able to process any refill requests and other follow-up actions by initiating text messages directly to patients. They were provided with a list each week containing action buckets such as “refill requests,” “change requests,” “date of birth authentication failed or incomplete,” “help requests,” “concerns about side effects,” and “other free text responses” and prioritized their handling of these action items. Figure 2 provides a view of the Engagement Console. Additional images of the Engagement Console are provided in Multimedia Appendix 1.

Initially, processing refill requests via the Engagement Console took an average of 10 to 15 minutes. After the first week, time needed to process refill requests via the Engagement Console dropped to about 5 to 10 minutes per patient.

## Results

### Refill Request Rate

Our primary process measure was the number of refill requests. Of 13,195 dialogues initiated, we received a total of 2405 text messages requesting refills (Table 2). These requests were then processed by the pharmacy team and tracked separately.


Table 3 shows the number of patients targeted and the percentage who refilled by patient list. The refill request rate was highest for the Near Goal patients (1581/8206, 19.27%).

**Table 2.** Refill request rate for text message group by month.

Month	Refill dialogues, n	Refill requests, n	Refill request rate, %
Month 1	6776	1140	16.82
Month 2	3190	647	20.28
Month 3	3229	618	19.14
3-month total	13,195	2405	18.23

**Table 3.** Refill request rate for text message group by adherence level.

Adherence level	Refill dialogues, n	Refill requests, n	Refill request rate, %
Near Goal	8206	1592	19.40
New Start	748	120	16.04
Nonadherent	4241	693	16.34
Group total	13,195	2405	18.23

**Figure 3.** Refills requests by hour from initial reminder.

### Time to Request Refill

Of those who requested a refill, 37.33% (898/2405) did so within 2 hours of receiving the initial reminder, an additional 48.61% (1169/2405) refilled within 24 hours (after also receiving the 2-hour reminder), and the remaining 14.05% (338/2405) refilled after receiving the 24-hour reminder. As displayed in Figure 3, there are spikes in refill activity immediately after the initial message (0), after the 2-hour reminder (2), and the 24-hour reminder (24). On average, members engaged within 24 minutes of getting the first message, and the median time to move through the refill process after engaging was less than 2 minutes.

Refill reminder dialogues were initiated between 10 am and noon on Wednesdays and Thursdays to allow for a reasonable time frame within which patients could respond. The bulk of refill requests (2210/2405, 91.89%) were made between the


hours of 10 am and 6 pm (Figure 4). A majority of responses were received within the first 4 hours, and 81.12% (1951/2405) of responses were received within the first 8 hours.

We tracked refill request processing by pharmacy staff (total of 8 KP staff members) and noted that they collectively processed about 40 to 50 refills in an hour by the end of the first month of the program. Anecdotal feedback from KP pharmacy staff suggests that this improvement in processing refill requests has allowed them to double monthly refills.

### Refills Processed

Our primary outcome measure was the number of refills that could be attributed to the text messaging. We were measuring the incremental effect of text messages (in addition to usual care) in increasing medication refills.



**Figure 4.** Percentage of refills requests by time of day.**Table 4.** Differences in refill rates between the text message and nontext message groups.

Month	Text message group refill rate, %	Nontext message group refill rate, %	Difference in refill rates Percentage points	P value
Month 1	35.73	23.49	12.24	.001
Month 2	52.55	39.10	13.45	.001
Month 3	54.05	43.23	10.82	.001
3-month total	44.08	30.01	14.07	.001

In the text message group, 12,272 patients received refill reminders via text (in addition to other outreach) over the 3-month program, and 5410 (44.08%) of these patients refilled their medication. The nontext message group of 76,068 patients received flyers and other outreach but no text reminders, and 22,826 (30.01%) of these patients completed medication refills (Table 4). The text message group refill rates were much higher than the nontext message group rates, and the 14.07 percentage point difference in refill rates between the 2 groups was statistically significant ( $P < .001$ ).

### Opt-Out Rates


The opt-out rate can be calculated in multiple ways and ranges from 1.02% to 5.09% depending on the calculation used. A total of 505 patients opted out over the course of the 3-month program. We have provided 3 different calculations in Table 5.

Here are the 3 different methods for calculating opt-out rates and rationale for each:

**Table 5.** Opt-out rates.

Approach for calculating opt-outs	Basis, n	Opt-out rate, %
Message level, messages	49,590	1.02
Dialogue level, dialogues	13,195	3.83
Member level, patients	9920	5.09

- Message level: This opt-out metric is calculated by dividing the number of members who opted out by the number of messages all members received. This measure helps us understand how long a member has stayed based on total volume of messages.
- Dialogue level: This opt-out metric is calculated by dividing the number of members who opted out by the number of dialogues all members received. This looks at the entire engagement in order to understand how well members received the program.
- Member level: This is the most common opt-out metric and is simply defined by dividing the number of members who opted out by the number of members at the beginning of the program. While this metric is useful, it does not factor in either program length or message volume and therefore presents a more coarse-grained view of member engagement and program value.

**Figure 5.** Sentiment in patient responses.

### Measuring User Experience

We analyzed patient free text responses to understand their experience and be more responsive. To do this, we used natural language processing to extract polarity, valence, and sentiment (very positive, positive, neutral, negative, very negative). For example, "Leave me alone" has a very different emotional tone than "Thanks so much for the reminder!" As shown in Figure 5, the largest subgroup of responses was neutral (1812/3609, 50.21%), followed by positive (1057/3609, 29.28%), very positive (434/3609, 12.03%), negative (301/3609, 8.34%), and very negative (5/3609, 0.14%).

### Ease of Use Survey Results

Another way in which we captured user experience was by asking patients directly. Starting in month 2, when patients completed a refill request, they received a confirmation message and were asked "Was this refill process easy to use?"

This question was intended to measure whether the TAM model's "ease of use" consideration had been successfully embedded in the refill dialogue solution. In designing for usability, we had prioritized the importance of creating a text-based refill dialogue that was easy to use, easy to learn, did not cause users to generate many errors, and was helpful to users. Over 70.02% (890/1271) of those who were presented with the survey question completed it. Of the 890 unique patients who completed the survey, 850 (95.51%) responded "Yes," and 40 (4.49%) responded "No."

## Discussion

### Principal Findings

We studied the value of an interactive text message refill solution with a chronically ill and partially adherent or nonadherent Medicare population and observed a difference of 14.07 percentage points in refill rates between the text message group and comparison group ( $P < .001$ ).

It is worth noting that patients in the texting group engaged at a much higher rate than predicted. We had estimated that the patient response rate would be between 10% and 20%, including stop requests, help requests, date of birth authentication attempts (successful and failed), refill requests, change requests, reasons for not refilling, and other free text responses. Our target refill request rate was 5% to 7% since we were messaging an older patient population. At the same time, we hoped that the ease of use of the refill dialogue might draw in more patients and nudge them toward completing their refill requests.

The program results far exceeded our expectations. Throughout the 3-month program, the response rate was around 37%, and the 3-month average refill request rate was 18%. We had also expected that since this was an older patient population the response time span might be stretched out a little longer, but this was not the case with over 80% of refill requests received within 8 hours of the initial reminder.

We used rules and basic natural language processing to improve recognition and handling of member responses over the course of the program, cut down unprocessed free text responses from



26% to under 16%, and reduced manual handling by pharmacy staff.

Overall patient feedback was very favorable and sentiment analysis of the responses revealed that patients were 5 times more likely to express positive sentiment than negative sentiment. Finally, almost 96% of the patients who completed refills via text message indicated that the solution was easy to use, and this strongly validated the TAM model and usability considerations that guided our design of the refill dialogues.

Although a cost-effectiveness analysis was not performed, interactive text messaging is inexpensive compared to manual calls or robo-calls. Finally, the high response rates and highly positive sentiment indicates improved patient engagement with their health care provider.

### Future Considerations

Our program incorporated basic demographic and psychographic data but did not tailor workflows based on the social determinants of health (ie, the conditions where people live, learn, work, and play and how these conditions affect their health risks and outcomes). This is an approach we plan to expand and implement in future programs. For example, how does living in a remote or rural area with no transportation impact refill behavior? How is income associated with refill rates? What about language and cultural barriers? This was a racially and ethnically diverse patient population. While the 3-month program used only English dialogues, the next phase would explore whether Spanish-speaking patients should be targeted differently and should also consider cultural and language barriers. We would also like to tailor content based on health literacy levels.

In future programs, we hope to combine demographic data (zip code, gender, age) with psychographic measures (adherence levels, past refill behavior, barriers to adherence, self-efficacy, stage of change, health beliefs) to develop a deeper understanding of the population being targeted to uncover health disparities and drive positive and sustained behavior change.

As we expand the program to other Kaiser Permanente regions, we expect to rely more heavily on machine learning-based natural language processing to improve recognition accuracy. Our machine learning-based natural language processing classifies a member's response into most commonly occurring categories which, in turn, triggers appropriate actions. We use a model that is topic-specific and trained on data that is based on a combination of responses received within the program and gathered through other sources. While we also rely on human intelligence to validate and handle outliers and unexpected responses, our goal is to reduce manual processing of member queries and responses to less than 5% in future programs.

### Conclusion

Findings suggest that partially adherent or nonadherent Medicare patients who receive interactive text message refill reminders have significantly higher medication refill rates compared to similar patients who do not receive text message refill reminders. The program results demonstrate that this incremental value of interactive text messages increased refill rates by 14.07 percentage points in Medicare patients.

Results of the program include increased refill rates and high levels of patient engagement. These results should provide insights for developing similar models that represent an elevated standard of care within patient management.

---

### Acknowledgments

We would like to thank the following individuals for their contributions to this project: Nico Arcino and Joyce Pedersen for initial development of project concept and scope; Allen Sarkisyan, Dustin Lo, and Ali Anari for design and development of data translation, loading, and presentation interfaces; Justine Espinoza and Nikta Moradi for overall project support; Chao Meng and Tony Askar for writing the underlying code for the dialogue engine within the mPulse Mobile platform; Matthew Betz for review and editing; and Paige Mantel for proofreading the manuscript.

---

### Conflicts of Interest

RBP, EF, MK, and RSP are current employees of mPulse Mobile, Inc, which is a vendor for Kaiser Permanente.

---

### Multimedia Appendix 1

Automated dialogues.

[[PDF File \(Adobe PDF File\), 2MB - mhealth\\_v6i1e30\\_app1.pdf](#)]

---

### References

1. Osterberg L, Blaschke T. Adherence to medication. *N Engl J Med* 2005 Aug 4;353(5):487-497. [doi: [10.1056/NEJMra050100](#)] [Medline: [16079372](#)]
2. Mahoney J, Ansell B, Fleming W, Butterworth S. The unhidden cost of noncompliance. *J Manag Care Pharm* 2008;14:S1-S29.
3. Iuga AO, McGuire MJ. Adherence and health care costs. *Risk Manag Healthc Policy* 2014;7:35-44 [[FREE Full text](#)] [doi: [10.2147/RMHP.S19801](#)] [Medline: [24591853](#)]

4. Viswanathan M, Golin CE, Jones CD, Ashok M, Blalock SJ, Wines RCM, et al. Interventions to improve adherence to self-administered medications for chronic diseases in the United States: a systematic review. *Ann Intern Med* 2012 Dec 4;157(11):785-795. [doi: [10.7326/0003-4819-157-11-201212040-00538](https://doi.org/10.7326/0003-4819-157-11-201212040-00538)] [Medline: [22964778](#)]
5. Latif S, McNicoll L. Medication and non-adherence in the older adult. *Med Health R I* 2009 Dec;92(12):418-419. [Medline: [20066833](#)]
6. Leslie RS, Tirado B, Patel BV, Rein PJ. Evaluation of an integrated adherence program aimed to increase Medicare Part D star rating measures. *J Manag Care Spec Pharm* 2014 Dec;20(12):1193-1203 [FREE Full text] [doi: [10.18553/jmcp.2014.20.12.1193](https://doi.org/10.18553/jmcp.2014.20.12.1193)] [Medline: [25443513](#)]
7. Levine DM, Lipsitz SR, Linder JA. Trends in seniors' use of digital health technology in the United States, 2011-2014. *JAMA* 2016 Aug 2;316(5):538-540. [doi: [10.1001/jama.2016.9124](https://doi.org/10.1001/jama.2016.9124)] [Medline: [27483069](#)]
8. Dailey J. Consumer survey on patient engagement.: Accenture; 2013. URL: [https://www.accenture.com/us-en/~media/Accenture/Conversion-Assets/DotCom/Documents/Global/PDF/Industries\\_11/Accenture-Silver-Surfers-Catching-eHealth-Wave-Infographic](https://www.accenture.com/us-en/~media/Accenture/Conversion-Assets/DotCom/Documents/Global/PDF/Industries_11/Accenture-Silver-Surfers-Catching-eHealth-Wave-Infographic) [accessed 2018-01-16] [WebCite Cache ID 6wWuXLF0B]
9. Anderson M. Technology device ownership: 2015. Washington: Pew Internet and American Life Project URL: [http://assets.pewresearch.org/wp-content/uploads/sites/14/2015/10/PI\\_2015-10-29\\_device-ownership\\_FINAL.pdf](http://assets.pewresearch.org/wp-content/uploads/sites/14/2015/10/PI_2015-10-29_device-ownership_FINAL.pdf) [accessed 2018-01-16] [WebCite Cache ID 6wWv6WkR1]
10. Rainie L. Digital divides. Washington: Pew Research Center URL: <http://www.pewinternet.org/2016/07/14/digital-divides-2016/> [accessed 2018-01-16] [WebCite Cache ID 6wWv2CJrr]
11. Thakkar J, Kurup R, Laba T, Santo K, Thiagalingam A, Rodgers A, et al. Mobile telephone text messaging for medication adherence in chronic disease: a meta-analysis. *JAMA Intern Med* 2016 Mar;176(3):340-349. [doi: [10.1001/jamainternmed.2015.7667](https://doi.org/10.1001/jamainternmed.2015.7667)] [Medline: [26831740](#)]
12. Pandey A, Krumme A, Patel T, Choudhry N. The impact of text messaging on medication adherence and exercise among postmyocardial infarction patients: randomized controlled pilot trial. *JMIR Mhealth Uhealth* 2017 Aug 03;5(8):e110 [FREE Full text] [doi: [10.2196/mhealth.7144](https://doi.org/10.2196/mhealth.7144)] [Medline: [28778843](#)]
13. Mira JJ, Navarro I, Botella F, Borrás F, Nuño-Solinis R, Orozco D, et al. A Spanish pillbox app for elderly patients taking multiple medications: randomized controlled trial. *J Med Internet Res* 2014;16(4):e99 [FREE Full text] [doi: [10.2196/jmir.3269](https://doi.org/10.2196/jmir.3269)] [Medline: [24705022](#)]
14. Ferreira F, Almeida N, Rosa AF, Oliveira A, Casimiro J, Silva S, et al. Elderly centered design for interaction and the case of the S4S medication assistant. *Procedia Computer Science* 2014;27:398-408. [doi: [10.1016/j.procs.2014.02.044](https://doi.org/10.1016/j.procs.2014.02.044)]
15. Dalgaard L, Gronvall E, Verdezoto N. MediFrame: a tablet application to plan, inform, remind and sustain older adults' medication intake. 2013 Presented at: Proceedings of the IEEE International Conference on Healthcare Informatics (ICHI); 2013; Philadelphia p. 36-45.
16. Silva J, Mouttham A, El Saddik A. UbiMeds: a mobile application to improve accessibility and support medication adherence. 2009 Presented at: Proceedings of the 1st ACM SIGMM International Workshop on Media Studies and Implementations that Help Improving Access to Disabled Users; 2009; Beijing p. 71-78. [doi: [10.1145/1631097.1631109](https://doi.org/10.1145/1631097.1631109)]
17. Vollmer WM, Owen-Smith AA, Tom JO, Laws R, Ditmer DG, Smith DH, et al. Improving adherence to cardiovascular disease medications with information technology. *Am J Manag Care* 2014 Nov;20(11 Spec No. 17):SP502-SP510 [FREE Full text] [Medline: [25811824](#)]
18. Kampmeijer R, Pavlova M, Tambor M, Golinowska S, Groot W. The use of e-health and m-health tools in health promotion and primary prevention among older adults: a systematic literature review. *BMC Health Serv Res* 2016 Sep 05;16 Suppl 5:290 [FREE Full text] [doi: [10.1186/s12913-016-1522-3](https://doi.org/10.1186/s12913-016-1522-3)] [Medline: [27608677](#)]
19. Penno E, Gauld R. Change, connectivity, and challenge: exploring the role of health technology in shaping health care for aging populations in Asia Pacific. *Health Systems & Reform* 2017 Aug 08;3(3):224-235. [doi: [10.1080/23288604.2017.1340927](https://doi.org/10.1080/23288604.2017.1340927)]
20. Davis FD. Perceived usefulness, perceived ease of use, and user acceptance of information technology. *MIS Quarterly* 1989 Sep;13(3):319-340. [doi: [10.2307/249008](https://doi.org/10.2307/249008)]
21. King WR, He J. A meta-analysis of the technology acceptance model. *Information & Management* 2006 Sep;43(6):740-755. [doi: [10.1016/j.im.2006.05.003](https://doi.org/10.1016/j.im.2006.05.003)]
22. Day M, Demiris G, Oliver DP, Courtney K, Hensel B. Exploring underutilization of videophones in hospice settings. *Telemed J E Health* 2007 Feb;13(1):25-31. [doi: [10.1089/tmj.2006.0023](https://doi.org/10.1089/tmj.2006.0023)] [Medline: [17309351](#)]

## Abbreviations

**CMS:** Centers for Medicare and Medicaid Services  
**DSR:** day supply remaining  
**EMR:** electronic medical records  
**KP:** Kaiser Permanente Southern California  
**PDC:** proportion of days covered  
**SMS:** short message service

**TAM:** technology acceptance model

*Edited by G Eysenbach; submitted 08.09.17; peer-reviewed by J Thakkar, J Redfern; comments to author 29.09.17; revised version received 20.10.17; accepted 08.01.18; published 30.01.18*

*Please cite as:*

*Brar Prayaga R, Jeong EW, Feger E, Noble HK, Kmiec M, Prayaga RS*

*Improving Refill Adherence in Medicare Patients With Tailored and Interactive Mobile Text Messaging: Pilot Study*

*JMIR Mhealth Uhealth 2018;6(1):e30*

URL: <http://mhealth.jmir.org/2018/1/e30/>


doi:[10.2196/mhealth.8930](https://doi.org/10.2196/mhealth.8930)

PMID:

©Rena Brar Prayaga, Erwin W Jeong, Erin Feger, Harmony K Noble, Magdalen Kmiec, Ram S Prayaga. Originally published in JMIR Mhealth and Uhealth (<http://mhealth.jmir.org>), 30.01.2018. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR mhealth and uhealth, is properly cited. The complete bibliographic information, a link to the original publication on <http://mhealth.jmir.org/>, as well as this copyright and license information must be included.



*This study is part one of a two-part study conducted over two years.  
Please continue reading for part two, released November 2019.*



# Impact of Social Determinants of Health and Demographics on Refill Requests by Medicare Patients Using a Conversational AI Text Messaging Solution: Cross-Sectional Study

AUTHORED BY



**KAISER  
PERMANENTE®**



PUBLISHED BY

**JMIR mHealth  
and uHealth**





Original Paper

# Impact of Social Determinants of Health and Demographics on Refill Requests by Medicare Patients Using a Conversational Artificial Intelligence Text Messaging Solution: Cross-Sectional Study

Rena Brar Prayaga<sup>1</sup>, MA, JD; Ridhika Agrawal<sup>1,2</sup>; Benjamin Nguyen<sup>1,2</sup>, BA; Erwin W Jeong<sup>3</sup>, PharmD; Harmony K Noble<sup>4</sup>, MSHCA; Andrew Paster<sup>1</sup>, BS; Ram S Prayaga<sup>1</sup>, MS

<sup>1</sup>mPulse Mobile, Inc, Encino, CA, United States

<sup>2</sup>Grinnell College, Grinnell, IA, United States

<sup>3</sup>Medicare Medication Therapy Management & Medication Adherence programs, Kaiser Permanente Southern California, Downey, CA, United States

<sup>4</sup>Regional Pharmacy Clinical Operations, Kaiser Permanente Southern California, Downey, CA, United States

## Corresponding Author:

Rena Brar Prayaga, MA, JD

mPulse Mobile, Inc

16530 Ventura Blvd

Suite 500

Encino, CA, 91436

United States

Phone: 1 888 678 5735

Email: [rena@mpulsemobile.com](mailto:rena@mpulsemobile.com)

## Abstract

**Background:** Nonadherence among patients with chronic disease continues to be a significant concern, and the use of text message refill reminders has been effective in improving adherence. However, questions remain about how differences in patient characteristics and demographics might influence the likelihood of refill using this channel.

**Objective:** The aim of this study was to evaluate the efficacy of an SMS-based refill reminder solution using conversational artificial intelligence (AI; an automated system that mimics human conversations) with a large Medicare patient population and to explore the association and impact of patient demographics (age, gender, race/ethnicity, language) and social determinants of health on successful engagement with the solution to improve refill adherence.

**Methods:** The study targeted 99,217 patients with chronic disease, median age of 71 years, for medication refill using the mPulse Mobile interactive SMS text messaging solution from December 2016 to February 2019. All patients were partially adherent or nonadherent Medicare Part D members of Kaiser Permanente, Southern California, a large integrated health plan. Patients received SMS reminders in English or Spanish and used simple numeric or text responses to validate their identity, view their medication, and complete a refill request. The refill requests were processed by Kaiser Permanente pharmacists and support staff, and refills were picked up at the pharmacy or mailed to patients. Descriptive statistics and predictive analytics were used to examine the patient population and their refill behavior. Qualitative text analysis was used to evaluate quality of conversational AI.

**Results:** Over the course of the study, 273,356 refill reminders requests were sent to 99,217 patients, resulting in 47,552 refill requests (17.40%). This was consistent with earlier pilot study findings. Of those who requested a refill, 54.81% (26,062/47,552) did so within 2 hours of the reminder. There was a strong inverse relationship ( $r=0.93$ ) between social determinants of health and refill requests. Spanish speakers (5149/48,156, 10.69%) had significantly lower refill request rates compared with English speakers (42,389/225,060, 18.83%;  $X^2_1 [n=273,216]=1829.2$ ;  $P<.001$ ). There were also significantly different rates of refill requests by age band ( $X^2_6 [n=268,793]=1460.3$ ;  $P<.001$ ), with younger patients requesting refills at a higher rate. Finally, the vast majority (284,598/307,484, 92.23%) of patient responses were handled using conversational AI.

**Conclusions:** Multiple factors impacted refill request rates, including a strong association between social determinants of health and refill rates. The findings suggest that higher refill requests are linked to language, race/ethnicity, age, and social determinants of health, and that English speakers, whites, those younger than 75 years, and those with lower social determinants of health



barriers are significantly more likely to request a refill via SMS. A neural network-based predictive model with an accuracy level of 78% was used to identify patients who might benefit from additional outreach to narrow identified gaps based on demographic and socioeconomic factors.

(*JMIR Mhealth Uhealth* 2019;7(11):e15771) doi: [10.2196/15771](https://doi.org/10.2196/15771)

## KEYWORDS

text messaging; SMS; refill adherence; medication adherence; Medicare patients; conversational AI; social determinants of health; predictive modeling; machine learning; health disparities

## Introduction

### Background

Medication nonadherence is when patients are unable to follow prescribed treatment dose, time of day, and frequency [1]. There are a range of factors, including patient-related, physician-related, and health system barriers, that contribute to nonadherence [2]. Adherence has been shown to have a significant effect on treatment outcomes [3,4] and has a major impact in managing chronic conditions such as hypertension, cardiovascular disease, and diabetes. The Centers for Medicare and Medicaid Services (CMS) define an adherent patient as someone whose proportion of days covered is greater or equal to 80%. Put another way, patients are considered adherent when they refill often enough to cover 80% or more of their medication plan as prescribed by their health care provider and as agreed to by the patient [5]. Dispensing or refill data is commonly used to compute adherence levels because of the validity, relative accessibility, and inexpensiveness of such data [6].

The World Health Organization estimates a medication nonadherence rate of 50% for patients with 1 or more chronic conditions [2,7]. This staggering proportion of nonadherence is estimated to annually cost between 100 to 290 billion dollars in the United States [8]. Moreover, nonadherence is estimated to cause approximately 125,000 deaths and at least 10% of hospitalizations every year [9,10].

### Medicare Patients and Adherence

Individuals suffering from multiple chronic conditions and taking multiple medications are more likely to be nonadherent [11]. Patients eligible for Medicare, who are individuals older than 65 years and/or who have a disability, fit this description of patients at greater risk of nonadherence. Older patients and those with disabilities have more chronic conditions and are usually on multiple medications. Of 586 Medicare recipients offered medication therapy management, 575 (98.1%) completed a survey that asked questions relating to adherence. Among those who responded, 406 (69.2%) reported that they took their medication regularly and as prescribed. Of the remaining 169 (30%), 123 identified forgetfulness as an issue, 18 (11%) mentioned side effects, and 17 (10%) said the medication was not needed. Lower adherence rates were associated with difficulty paying for medication. Finally, subsidy recipients and non-English speakers were significantly less likely to be counseled about drug side effects [12].

### Use of Mobile Technology for Adherence

In a 2018 press release, the CMS committed to supporting modern and virtual methods of health care [13]. Furthermore, in a multinational survey conducted in 2018, 77% of respondents said the ability to request prescription refills via text message would increase their likelihood of choosing a health care provider. This percent is a 10-point increase over a 3-year span [14]. The Deloitte Center for Health Solutions conducted a nationally representative survey in which approximately a third of individuals indicated interest in receiving text messages for nutrition, exercise, sleep, and stress management [15]. These trends represent a changing societal landscape, and the health care field is poised to address this identified need. A recent interactive mobile solution for appointment reminders within the Department of Veterans Affairs (VEText) has been used to send SMS text message reminders to over 6 million veterans [16,17]. Text messages can also provide links to resources and reminders toward adopting healthier behaviors. Several meta-analyses corroborate the effectiveness of SMS for medication adherence [18-20]. An earlier study by the authors [21] measured the impact of SMS text reminders on refill rates of nonadherent and partially adherent Medicare patients with chronic disease. They found that text reminders increased refill rates by 14 percentage points compared with those who did not receive these reminders.

Important to the success of any intervention is its implementation, scalability, and sustainability [22]. Text messaging presents an effective, affordable, and scalable tool [21] that can use conversational AI to greatly impact health outcomes. More specifically, conversational AI (or conversational agents) can encourage health care consumers to engage with systems that imitate human conversations using text [23,24]. For example, a fully automated conversational AI system has been used to promote weight loss among overweight and obese diabetic patients [25]. As conversational agents can learn over time, interventions with thousands of users can be used to inform and improve the quality of the conversations, often within days or weeks. However, there is limited research currently available on the use of conversational AI within SMS (and not app-based) messaging for refill adherence.

### Social Determinants of Health

The World Health Organization has defined social determinants of health (SDOH) as the conditions or circumstances in which people are born, grow, live, work, and age [26]. The most commonly identified SDOH in the United States are housing, income, food, transportation, education, race/ethnicity, and unemployment [27,28]. Despite notable improvements in overall

health over the past few decades, inequalities of SDOH contribute to persistent disparities in life expectancy and health outcomes [29,30].

Social determinants have also been linked to nonadherence, and a recent study using data from the National Health Interview Survey found that half of the adults with diabetes perceived financial stress, while one-fifth reported financial insecurity and food insecurity [30]. Since SDOH are not always recognized, they might be overlooked by a clinician in a medical setting. In a study of patients with chronic disease, two-thirds of those who reported not taking medications as prescribed due to cost never shared this with their physician [30,31]. The National Academy of Medicine has published a framework for educating health care professionals on the importance of SDOH [32], but there is still limited research studying the specific ways in which SDOH pathways interact to impact adherence, particularly in older populations.

## Objectives

A pilot study examining Medicare Part D patients over a period of 3 months supported the value of using SMS text message refill reminders to increase medication refill rates [21]. This study increases the sample size of those receiving SMS refill reminders to 99,217 Medicare recipients, includes both English and Spanish language text messages (the pilot used only English messages), and expands the duration to a 2-year follow-up period. This analysis focuses on a few different questions:

- First, were the results from the pilot study replicable with a much larger population using an enhanced version of the text messaging solution with improved conversational AI?
- Second, is there a relationship between SDOH and refill request rates, how large is this association, and do SDOH attenuate the association?
- Third, how do other variations in patient characteristics (age, gender, ethnicity, and language) moderate and predict

likelihood of requesting a refill using a text message solution?

## Methods

### Participants

The SMS refill reminder program began as a 3-month pilot in December 2016 and was expanded to include multiple regions within a large integrated health system. The analysis covers a 2-year period from December 2016 to February 2019. It includes a population of 99,217 English- and Spanish-speaking patients (median age 71 years) targeted for medication refill by Kaiser Permanente, Southern California. The Kaiser Permanente, Southern California, Institutional Review Board determined that this program did not require review and was exempt.

All patients had Medicare Part D as their pharmacy benefits and had one or more chronic conditions (diabetes, hypertension, high cholesterol, and/or anticoagulation). Patients in this program refilled one or more of the following 4 classes of drugs: oral diabetes medications, blood pressure medications (renin-angiotensin system antagonists), statins, and direct oral anticoagulants (DOAC).

Targeted patients were shared by Kaiser Permanente, Southern California, in a weekly file, and they had varying levels of nonadherence. The total number of patients targeted for refill ranged from 1000 to 9000 patients per month. Note that these patients were not distinct each month because they could be on the list several times in a year. Patient records included first name, date of birth (DOB), gender, spoken language, address, race/ethnicity, mobile phone number, opt-in status, and refill drug(s). These fields, when available, were used for all analysis.

Tables 1 and 2 provide age and race/ethnicity breakdowns for this group.

**Table 1.** Age of text messaging group.

Age band (years)	Patients, n (%)	Reminders, n (%)
<60	6344 (6.39)	20,883 (7.64)
60-65	4502 (4.54)	13,806 (5.05)
65-70	29,600 (29.83)	79,349 (29.03)
70-75	27,201 (27.42)	75,902 (27.77)
75-80	15,039 (15.16)	42,221 (15.44)
80-85	7899 (7.96)	22,247 (8.14)
>85	5458 (5.50)	14,385 (5.26)
Unspecified	3174 (3.20)	4563 (1.67)
Total	99,217 (100)	273,356 (100)

**Table 2.** Race/ethnicity of text messaging group.

Race/ethnicity	Patients, n (%)	Reminders, n (%)
White	30,683 (30.93)	81,544 (29.83)
Hispanic/Latino	21,841 (22.01)	67,266 (24.60)
Black/African American	9124 (9.20)	28,365 (10.38)
Asian	8705 (8.77)	23,870 (8.73)
Other/mixed	1372 (1.38)	3812 (1.39)
Unspecified/unknown	27,492 (28.71)	68,499 (25.06)
Total	99,217 (100)	273,356 (100)

## Procedure


The intervention used the mPulse Mobile platform to deliver SMS text messages to patients. Patients in the text messaging group received a refill reminder dialogue that consisted of a series of messages written at a sixth-grade readability level.

Text message refill reminders were sent out on a weekly basis to patients who were due for a refill. The first message was a greeting, reminding patients that they were due for a refill. They were then prompted to validate their DOB (to ensure the person was the intended recipient of the reminder) by choosing from one of five options. If the patient validated their DOB successfully, they could then view their medication(s) due for refill and the last filling pharmacy. Patients did not get a second chance if they did not validate correctly. In those cases, a member from the pharmacy staff would reach out to discuss barriers to nonadherence and/or complete the refill by phone. As part of the refill workflow, patients could select whether to receive their medication by mail (this was added as an option

in May 2018) or to pick up their medication(s) at a Kaiser Permanente outpatient pharmacy. They could also change their default pickup pharmacy. Pharmacy staff who were located at a central location were involved with managing the responses that came back from the patients. These patient responses were categorized as “Refill request,” “Barriers,” “Date of birth issue,” “Free text response,” “Side effects,” “Change Pharmacy,” and “Help.”

As in the pilot study [21], we used mPulse Mobile’s Engagement Console to support the pharmacy staff. This Web-based user interface allows users to quickly identify and prioritize subgroups of patients (as described above) for quicker follow-up.

Figure 1 provides a view of the refill reminder message flow and the various steps or options within this dialogue. The initial step reminds the patients to refill the medication(s) and requests that they respond with the *structured options* “1” to continue or “2” to end. However, in some instances, patients respond with *unstructured responses* such as “I’ve already refilled” or “No thanks.”

**Figure 1.** Overview of message flow within refill dialogue.

A patient would receive a maximum of 3 messages if they did not respond (initial reminder, 2-hour reminder, and 24-hour reminder). The patient could opt out at any point during the messaging flow, and no subsequent messages would be sent.

Conversational AI was developed and used within the solution to also automatically process the following types of unstructured responses and patient requests:

- Patient wants to unsubscribe from texting program but does not reply “STOP” or 7867 (patients with older phones could use number keys to correspond to letters).
- Patient confirms intent to request a refill by using an unexpected phrase.
- Patient is experiencing side effects and might require medical attention.
- Patient wants to change pharmacy location where they wish to pick up the refill.
- Patient does not want to refill and provides a reason for not refilling.
- Patient requests help or wants additional information.
- Patient provides correct DOB instead of selecting a numeric option.
- Patient wants to switch language (English to Spanish or vice-versa).

For purposes of the qualitative analysis, each response was strictly coded as structured or unstructured. A response was considered *structured* if it exactly matched any of the following strings: 0, 1, 2, 3, 4, 5, 6, 7867, and case insensitive versions of *AYUDA*, *HELP*, *MAIL*, *RESUB*, *STOP*, and *STOPALL*). All other responses were coded as unstructured.

Two coders classified each conversational AI rule using the description within the solution (eg, DOB validation and change order). There were a total of 75 conversational AI rules. These individual rules were then combined into 13 broader “rule type” categories: *initial reply*, *refill process*, *change pharmacy*, *DOB validation*, *barriers*, *payment*, *subscription*, *language change*, *member information*, *acknowledgment*, *feedback*, *help*, and *did not understand*. All rules belonged to the 13 rule types, and any ambiguity about rule type were resolved by discussion and agreement between the two coders.

An internally developed SDOH index was used to understand how unmet needs might impact patient refill behavior. When patient address was available, an SDOH index was computed. [Multimedia Appendix 1](#) outlines the factors that were used to compute the SDOH index and to create low, medium, and high SDOH clusters.

A neural network multilayer perceptron (MLP) model was used to perform predictive analysis on factors that might impact refill requests.

## Results


The results of the scaled intervention are summarized in 3 parts: (1) a replication of the analysis performed in the pilot study; (2) results of subgroup exploratory analysis, including the use of an SDOH index and a predictive model; and (3) a qualitative analysis of the use and value of conversational AI and interactivity.


### Part 1: Analysis of Scaled Program

A total of 273,356 SMS reminders were sent over a 2-year period to 99,217 Medicare Part D patients who had opted in to texting. In response, 17.40% (47,552/273,356) refills were requested. [Figure 2](#) shows the conversion funnel from reminder to refill request.

DOB validation was a necessary step to view the refill information, and this step resulted in a drop-off (DOB validation failures or did not attempt) of 6.55% (6288/95,121). Of those who requested a refill, 54.81% did so within 2 hours after receiving the initial reminder (N=26,062/47,552). As displayed in [Figure 3](#), there are spikes in refill activity immediately after the initial message (“0”), after the 2-hour reminder (“2”), and the 24-hour reminder (“24”).

**Figure 2.** Conversion of refill reminder to refill request. DOB: date of birth.



**Figure 3.** Refills requests by hour from initial reminder.

## Part 2: Exploratory Subgroup Analysis

We present the following subgroup analysis using 4 variables: SDOH, language, race/ethnicity, and age (gender did not have a significant moderating effect on refill rates).


### Social Determinants of Health Analysis

Patients were grouped into 10 evenly spaced SDOH bands from 0 to 100. Refill requests were very highly inversely correlated with SDOH bands ( $r = -0.93$ ), as shown in Figure 4.

To further understand the impact of SDOH on refill process, we grouped the SDOH bands further into 3 SDOH clusters


(high, medium, and low) using k-means clustering as described further in [Multimedia Appendix 1](#).

As can be seen in Table 3 and Figure 5, the negative correlation of refill request rates to SDOH index is driven primarily by the initial response rates (ie, after receiving the “Welcome” message in Figure 1, the patient confirms their intent to move forward in the dialogue). The difference in average SDOH between those who reply and do not reply was statistically significant ( $t_{252,834} = -55.07$ ;  $P < .001$ ), but there was no impact of SDOH for refill requests after the patient engages with the initial text message ( $t_{87,234} = 1.71$ ;  $P = .09$ ). In other words, once a patient is willing to engage with the texting program, they request refills at the same rate regardless of SDOH levels.

**Figure 4.** Refill rates versus social determinants of health bands.

**Table 3.** Refill request rate for text message group by social determinants of health level.

Social determinants of health	Refill dialogues, N	Percent who responded, n (%)	Percent of responders who requested refill, n (%)
Low (0-52.8)	124,423	47,949 (38.54)	23,623 (49.27)
Medium (52.8-86.1)	94,489	30,755 (32.55)	15,464 (50.28)
High (86.1-100)	33,924	8532 (25.15)	4357 (51.07)

**Figure 5.** Social determinants of health impact on response rates and percentage of responders who request refill. SDOH: social determinants of health.

### Spanish Versus English

Spanish-speaking patients had significantly lower refill request rates (5149/48,156, 10.69%) compared with English-speaking patients (42,389/225,060, 18.83%;  $X^2_1$ , [n=273,216]=1829.2;  $P<.001$ ). As with SDOH impact, the initial response rates also vary by spoken language, where Spanish-speaking patients (8984/48,156, 18.66%) were far less likely to engage with text messaging than English-speaking patients (86,105/225,060, 38.26%;  $X^2_1$ , [n=225,060]=6716.9;  $P<.001$ ).

Interestingly, a difference in refill request rates by language after the patient engaged with the reminder shows that Spanish-speaking patients request refills at a higher rate compared with English-speaking patients once they engage (5149/8984, 57.31% vs 42,386/86,105, 49.22%;  $X^2_1$  [n=95,089]=212.5;  $P<.001$ ).

We used a pointwise biserial correlation (point biserial correlation  $r_{pb}=0.27$ ;  $P<.001$ ;  $N=252,696$ ) to find that higher SDOH values are correlated with Spanish language preference. Spanish speakers ( $N=44,869$ ) had a higher average SDOH of 71.78 as compared with English speakers' ( $N=207,827$ ) average of 55.65 ( $t=151.39$ ;  $P<.001$ ).

### Age

Younger patients were significantly more likely to reply and request refills compared with older patients ( $t_{83,415}=-43.30$ ;  $P<.001$ ). The older age group (75 years and older) responded at a rate of 29.84%, whereas patients younger than 45 years responded at a rate of 47.81%. There were also significantly different rates of refill requests by age band ( $X^2_7$  [n=268,793]=1460.3;  $P<.001$ ), with younger patients requesting refills at a higher rate, as shown in Table 4. We do see a spike in refill requests in the 85+ years group, and this suggests that caregivers or family members might be more actively assisting patients in this age band.



**Table 4.** Response and refill request rates by age.

Age band (years)	Refill dialogues, N	Responded, n	Date of birth validation, n	Refills requested, n	Request rate, %
<60	20,883	9388	8929	5277	25.27
60-65	13,806	5444	5061	2808	20.34
65-70	79,349	29,625	27,787	15,014	18.92
70-75	75,902	25,707	23,949	12,458	16.41
75-80	42,221	127,732	11,781	6103	14.45
80-85	22,247	6256	5780	3039	13.66
>85	14,385	4538	4242	2351	16.34
Unspecified	4563	1431	1364	501	10.98
Total	273,356	95,121	88,893	47,552	17.40


### ***Does Race/Ethnicity Have an Influence on Engagement and Refill Request Rates?***

Patient race/ethnicity had a significant effect on initial response rates ( $X^2_4$  [n=204,857]=5282.40;  $P<.001$ ). Patients who identified as white responded at the highest rate (34,134/81,544,

41.86%), whereas Hispanic/Latino patients had the lowest response rate (16,700/67,266, 24.83%). Once someone did respond, race/ethnicity did not influence whether they refilled or not ( $X^2_4$  [n=68,329]=2.37;  $P=.07$ ), as reflected in Table 5 and Figure 6.

**Table 5.** Response and refill request rates by race/ethnicity.

Race/ethnicity	Refill dialogues, N	Responded, n	Date of birth validation, n	Refills requested, n	Request rate, %
Asian	23,870	6695	6233	3296	13.81
Hispanic/Latino	67,266	16,700	15,430	8711	12.95
Black/African American	28,365	9476	8751	4678	16.50
Other/mixed	3812	1324	1216	652	17.10
White	81,544	34,134	32,181	16,557	20.30

**Figure 6.** Impact of race/ethnicity on response rates and percentage of responders who request refill.

### ***Predictive Model to Improve Refill Adherence***


An in-depth analysis of the results revealed that the *reply rate* was the most important variable that we could influence to drive refill rates. After replying to the reminder, about half of the

patients continue the conversation to validate their DOB and then request a refill of their medication(s), while the other half drop off and do not engage further.

As we were uncertain of the interaction of age, gender, SDOH, language, and race/ethnicity and how they moderate reply rates,

we built a machine learning model using these factors as inputs to predict those least likely to reply. We trained a supervised model to predict reply likelihood based on known attributes. The model consisted of a neural network multilayer perceptron (MLP) with 24 input units, 10 hidden units, and a single output unit (to represent reply likelihood). The training set consisted of 70% of the available data, and model validation was done with the remaining 30%. The input vector consisted of features representing age, race/ethnicity, gender, SDOH, language, drug class, etc. We selected the most discriminative features to avoid overfitting and to reduce multicollinearity and redundancy in the feature space, and we used one-hot encoding to ensure uniform scale across features. Finally, we excluded rows with incomplete values. Each data row represented an instance of a unique series of events after a reminder was sent to a patient. This meant that if a member had been contacted 3 times to refill a drug and they only replied twice, the “reply” value would now be 0.66. This solved the problem of contradictory data and also converted the reply variable into a continuous variable representing reply likelihood.

**Figure 7.** Minimizing the number of false positives.



### Part 3: Qualitative Analysis of the Usefulness of Conversational Artificial Intelligence

Our solution incorporated conversational AI (CAI) and natural language understanding (NLU) to provide a robust and successful interactive experience that ensures that the patient is able to request their refill as quickly and conveniently as possible.

We performed a qualitative analysis of all patient responses to evaluate whether the more complex capabilities of CAI and NLU were necessary and helpful for a better patient experience. As part of this analysis, we coded 307,484 responses as either structured or unstructured (as described in the Methods section).

Of the 307,484 responses that we received during the study period, only 7.77% (n=23,886) were not understood by the CAI. Table 6 provides a breakdown of structured and unstructured responses as well as the steps in the refill reminder dialogue

### Optimizing the Model to Predict Those Least Likely to Engage and Request Refill

To develop an approach to impact refill adherence, we wanted to first maximize the prediction accuracy for patients who were not likely to reply at all. We were less concerned about prediction accuracy for patients who were likely to engage and request refills. For the model to have value in an applied setting, we wanted to capture as many patients who were not likely to engage and might require additional support to request a refill. Figure 7 contains a visual representation of the confusion matrix, which highlights only those predictions that relate to nonreplying members.

To address these dual objectives (accurately predicting those who require outreach while maximizing the number of people who require outreach), we found a cut-off of 37% predicted likelihood of replying as a good threshold. This means that a patient whose predicted likelihood is less than 37% should fall into a category of requiring additional outreach. In summary, we can identify over 66% of those requiring outreach (because they will not reply) at a model accuracy of 78%.

where they typically occurred. All unstructured responses shown in Table 6 were understood by the CAI engine, which triggered the appropriate replies.

There were several instances when asked to provide structured response (eg, text 1 to view the medication), a member replies with additional information (eg, “1 – I only took this medication for 1 day and it caused great muscle pain. In the meantime, my cholesterol levels are now below 200”). Similarly, when asked to validate DOB by choosing from 1 of the 3 options, members will choose a number but also input the DOB as in “1 - 1950-05-30.” The system was largely successful in recognizing responses and accurately categorizing them. In Table 7, we present additional examples where the CAI was successful.

Textbox 1 includes a few sample responses that we failed to understand but could have handled appropriately if the patient had responded when expected or provided more context. In summary, our results indicate that despite the overall high

accuracy (over 92%) of handling patient responses by the CAI, there are a number of instances (almost 8%) that were not handled at all, and in these cases, the patient was informed that the system was unable to understand their message.

**Table 6.** Counts of the type of responses that were handled by the conversational artificial intelligence (CAI).

Role of CAI	Structured responses	Unstructured responses	Total, n (%)
Supported by CAI	272,364	11,234	284,598 (92.23)
CAI could not handle	3001	20,885	23,886 (7.77)
Total	275,365 (89.55)	32,119 (10.45)	307,484 (100)

**Table 7.** Examples of unstructured patient messages when the conversational artificial intelligence successfully understood the message.

Sample patient responses	Response category
"1-I only took this medication for 1 day and it caused great muscle pain. In the meantime my cholesterol levels are now below 200"	Side effect barrier
"1/2 tablet twice or thrice weekly. Changed on 6/25/18 due to recurring muscle pain and Dr XXXX concurred"	Side effect barrier
"Both kinds of pain. I can't walk very far if i take too many so I've cut it to 1/4th and then when it gets to bad i stop it for a few days."	Side effect barrier
"Caused pain on calvrs so bad i could not walk"	Side effect barrier
"Doctor took me off medication because of too much muscle pain"	Side effect barrier
"I cant take a Statin It gives me terrible muscle pain and I cant sleep !!!! No."	Side effect barrier
"I have reported to my previous primary and his medical assistant...that the dosage prescribed gave me leg cramps and pain...I agreed to take half a pi"	Side effect barrier
"Do not have the money right now to fill it will refill it a when I have the funds"	Cost barrier
"Don't have money right now"	Cost barrier
"Don't have the extra money till the first."	Cost barrier
"Don't have the money yet to refill but plan on refilling soon"	Cost barrier
"Filling px on the military base at no cost"	Cost barrier
"1 (one)"	"1"
"1 (sent with Invisible Ink)"	"1"
"1 proceed to refill"	"1"
"1 refill"	"1"
"1 -"	"1"

**Textbox 1.** Sample of unstructured patient messages when the conversational AI responded to patients that it did not understand their message.

Sample patient responses

“0 - This has already been ordered last weekend!”

“2 cuando lo recojo?”

“2 my body can’t tolerate this drug”

“A refill is not needed until the end of march.”

“Already ordered online”

“Cancel rx change”

“Doctor has lowered dosage from prescribed amount. Doctor needs to update prescription.”

“I am in San Diego now”

“I am out of the country”

“I need talk to Sam body”

“I want to talk to my doctor first.”

“Random”

“Still have full bottle of 60. Will order later.”

“What medication??”

## Discussion

### Principal Findings

We found that the refill request rates reported in the pilot study with English speakers [21] could be replicated at scale. The results in this study confirm and replicate the pilot results and reveal that, even after significant expansion of the population (scaled to over 7 times the initial group size and with the addition of Spanish speakers), the solution was effective in moving patients through the text dialogue to quickly complete a refill request. Pilot refill request rates of 18.1% were closely mirrored (17.40%), or even higher when limited to requests by English speakers (18.83%). Expanding the capabilities to include Spanish speakers was an important feature that is not commonly available in other texting solutions.

On the basis of the results in the previous section, we found that patients are not equally likely to request a refill. The findings suggest that good refill adherence is linked to language, race/ethnicity, age, and SDOH and that English speakers, white patients, those younger than 75 years, and those with lower SDOH barriers have significantly higher odds of requesting a refill via SMS. We studied these associations and developed a tool and approach that can be used for future outreach to narrow identified gaps based on demographic and socioeconomic factors and to increase overall refill adherence.

Finally, we wanted to evaluate the impact of the conversational AI engine, which is not typically found in other health care texting solutions. The results indicate that patients needed conversational AI as they traversed the refill reminder dialogue, and as health care consumers become more comfortable and familiar with artificial intelligence-based agents and chatbots, this expectation will only increase.

To our knowledge, no other published study (other than the study by Brar Prayaga et al [21]) has reported a refill adherence

solution with results at scale [20]. The closest comparable solution with a large volume of patients is the VEText system for appointment reminders offered by the Department of Veterans Affairs, and we look forward to an in-depth study of that solution.

### Understanding Who Engages and Requests a Medication Refill

As our results indicate, we have identified several associations between factors that impact a patient’s likelihood to reply to the reminder and continue on to request a refill via SMS text messaging. These include language, SDOH, age, and race/ethnicity. For example, Spanish speakers were much less likely to engage with the reminder at all and, therefore, requested refills at a significantly lower rate of 10.69%, compared with English speakers at 18.83%. Note that for purposes of this text message solution, we messaged all patients in English (including those with a preferred language of Tagalog, Mandarin, etc in their health record) unless they had a preferred language of Spanish. Patients who identified as Hispanic/Latino had the lowest refill request rates of 12.95%, followed by Asian (13.81%), black (16.50%), and white (20.30%). Interestingly, neighborhood-based SDOH levels were highly correlated with patient language (English/Spanish), and patients with high SDOH barriers had significantly lower refill request rates of 12.84%, compared with medium SDOH (16.37%) and low SDOH levels (18.99%), as shown in Figure 5. Finally, younger patients were significantly more likely to reply and request refills compared with older patients. We believe these associations and effects should be addressed and have developed a predictive tool to help improve overall refill adherence rates for Medicare D patients.

### Using a Predictive Model to Assign Resources

A key goal is to increase the initial reply rate, and we were most interested in uncovering the population (roughly two-thirds) who were *unlikely to engage at all*. We found that the predictive

model is able to accurately (>78%) pinpoint a high percentage (>66%) of patients who will not engage with an SMS text reminder to request a refill. This can be used as a valuable tool by a health provider or pharmacy to proactively communicate with populations who are least likely to complete a refill request. While current outreach methods to encourage adherence, such as phone calls by pharmacy staff or automated interactive voice response calls, are typically more costly and time consuming [21,33], a targeted approach using the predictive model could optimize limited staff resources.

### Addressing Barriers to Improve Refill Adherence

Another recommendation is to reduce the *initial barriers* or unwillingness to engage with the program. For example, lack of familiarity with texting or mistrust of the channel, especially among older patients, could be addressed with more tailored versions of the initial reminder that alleviate possible unease with using SMS text messaging to complete a refill transaction.

Similarly, health plans and providers could provide supplemental resources (such as an informational video to explain the process for requesting a refill via text message and to address any specific concerns of Spanish speakers), and the text dialogue could link to these resources. It might also be beneficial to add multilanguage support to expand beyond English and Spanish. In addition, as cost can be a barrier to refill for patients, especially for those with high SDOH levels, the text message dialogue could remind patients that, for instance, mail order refills are incentivized (eg, “did you know that if you order by mail you can get a 3-month supply of medication for the cost of a 2-month supply?”).

We also found that for a large percentage of refill reminders (15.75%, n=14,005), patients start the refill process (ie, view medication and validate their DOB) and then share a barrier or other concern that causes them to drop out of the process. There were an additional 10,192 instances where patients shared a reason for not refilling in other parts of the conversational flow. These barriers include a wide range of topics such as cost, side effects, already refilled, still have sufficient medication, do not want to refill, and taking differently than prescribed. We are sharing this response data with Kaiser Permanente, Southern California, and they are continuing to find ways to address these issues and follow-up with patients as required.

### The Role of Conversational Artificial Intelligence

Conversational AI was helpful in moving patients through the refill dialogue using mostly structured inputs almost 90% of the time. Although complex conversational AI was not essential for driving outcomes (validating DOB, completing refill, etc), it played an important role in keeping the conversation going when patients engaged using unprompted or unstructured messages (“Caused pain on calvrs so bad I could not walk [sic]”). In this example, it is recognizing the patient’s expressed concern as a *side effect*, thereby allowing the member to respond as they see fit to continue the conversation instead of being constrained by a rigid structure. This model of primarily relying on prompted responses and also understanding those cases where users want to go outside a closed response set allows the user to control the flow of the conversation in keeping with Grice’s

Maxim of Manner [34]. Press releases relating to the recent launch of VEText [16] suggest a significant impact on appointment no-show rates at scale (there is no peer-reviewed study on the solution currently available, but the press release reported a no-show reduction from 13.7% to 11.7%, N not reported). The system requires users to only use prompted and structured alphanumeric responses, such as R2 and K3 [17]. While the details of the VEText solution are unclear, support for unstructured responses (estimating 10% based on our results) using conversational AI would likely improve user experience as well as overall outcomes. We believe this hybrid approach of supporting dual modes of interaction will support a fluid and frictionless conversation to enable task completion and allows a more empathetic exchange with the health care partner. Finally, while our accuracy rate of conversational AI was over 92%, we continue to explore ways to improve the system. A review of the literature using conversational AI reveals that the focus area is limited in scale and tends to be restricted to apps [24] and not SMS.

It is unclear why there is limited adoption of conversational AI within SMS text messaging as this is a channel with potential to reach all segments of the population. This study is unique in analyzing a text messaging and conversational AI solution at scale that allows elderly populations to easily and conveniently request their medication refills.

### Limitations

The findings of this study have to be seen in light of some limitations. The study was not a randomized controlled trial, and there is the possibility of selection bias. Due to regulations within the wireless communication industry and the Telephone Consumer Protection Act, we must have prior consent before messaging patients, and this constraint applies to any automated text messaging solution. As a result, the study targeted only those patients who had already opted into digital engagement. As the messaging program requires that patients have a mobile number with a texting plan, patients with only landline numbers were excluded. The nontext group received phone call outreach as part of the standard of care. However, we have already demonstrated the value and incremental benefit of SMS text reminders as compared with phone reminders [21], and this was not a focus of the study.

Using bivariate analyses can inflate the type I error, and this is a limitation. At the same time, we attempted to address this issue by using a neural network predictive model, which is a form of multivariate regression and can reduce the impact of multicollinearity. In addition, we did not address the impact of multiple reminders over time—that is, does a patient who requests a refill after receiving a reminder, and later becomes nonadherent again, continue this positive behavior upon receiving a future reminder?

Finally, the solution described in this study is a commercial system offered by mPulse Mobile with a licensing fee. We are unable to share financial benefits of using the system, such as operational efficiencies, health savings, and reimbursement revenue from improved adherence, as this information is proprietary and confidential. At the same time, we cannot disclose solution costs (such as messaging costs, license fees,



and implementation costs). However, the overall financial gains from using the refill reminder solution were greater than system costs.

## Conclusions

Overall, this study indicates that there are sharp differences in likelihood to reply to a refill reminder and complete a refill request via SMS based on demographic and socioeconomic factors. We found a strong association between refill request rates and patient language, age, race/ethnicity, and SDOH levels, and these differences may contribute to health disparities and

impact health outcomes in Medicare patients. Using a predictive and innovative model to target patients least likely to engage with the SMS solution and crafting a tailored mobile communication and conversational AI strategy could reduce these inequalities and improve refill adherence. We will continue to refine our solution and optimize our predictive model to validate our results and hope to be able to address disparities and drive even stronger outcomes. Finally, we believe that, to ensure the success of a text messaging solution and yield similar results, message tone and content, ease of use, level of tailoring, and quality of conversational AI are important considerations.

## Acknowledgments

The authors would like to thank the following individuals for their contributions to this project: Joyce Lin for data translation; Bhumika Gupta for support with conversational AI; Emily Haag for review of member responses; Kayvon Moradi, Michael Steinmetz, and Heather Forst for overall project support; and Carlton Segbefia for help with code optimization and analysis.

## Conflicts of Interest

RBP, AP, and RSP are current employees of mPulse Mobile, Inc, which is a vendor for Kaiser Permanente. RA and BN contributed to this study as part of their internship at mPulse Mobile, Inc.

## Multimedia Appendix 1

Calculating the Social Determinants of Health Index.

[[PDF File \(Adobe PDF File\), 93 KB-Multimedia Appendix 1](#)]

## References

1. Hugtenburg J, Timmers L, Elders PJ, Vervloet M, van Dijk L. Definitions, variants, and causes of nonadherence with medication: a challenge for tailored interventions. *Patient Prefer Adherence* 2013;7:675-682 [[FREE Full text](#)] [doi: [10.2147/PPA.S29549](#)] [Medline: [23874088](#)]
2. de Geest S, Sabaté E. Adherence to Long-term Therapies: Evidence for Action. Geneva, Switzerland: World Health Organization; 2003.
3. Modi AC, Quittner AL. Barriers to treatment adherence for children with cystic fibrosis and asthma: what gets in the way? *J Pediatr Psychol* 2006 Sep;31(8):846-858. [doi: [10.1093/jpepsy/jsj096](#)] [Medline: [16401680](#)]
4. DiMatteo MR, Haskard-Zolnier KB, Martin LR. Improving patient adherence: a three-factor model to guide practice. *Health Psychol Rev* 2012;6(1):74-91 [[FREE Full text](#)] [doi: [10.1080/17437199.2010.537592](#)]
5. Centers for Medicare & Medicaid Services. 2018. Medicare-Medicaid Plan Performance data Technical Notes. URL: [https://www.cms.gov/Medicare-Medicaid-Coordination/Medicare-and-Medicaid-Coordination/Medicare-Medicaid-Coordination-Office/FinancialAlignmentInitiative/Downloads/MMPPPerformanceDataTechNotesCY2018\\_04252018.pdf](https://www.cms.gov/Medicare-Medicaid-Coordination/Medicare-and-Medicaid-Coordination/Medicare-Medicaid-Coordination-Office/FinancialAlignmentInitiative/Downloads/MMPPPerformanceDataTechNotesCY2018_04252018.pdf) [accessed 2019-06-13]
6. Torres-Robles A, Wiecek E, Cutler R, Drake B, Benrimoj SI, Fernandez-Llimos F, et al. Using dispensing data to evaluate adherence implementation rates in community pharmacy. *Front Pharmacol* 2019;10:130 [[FREE Full text](#)] [doi: [10.3389/fphar.2019.00130](#)] [Medline: [30863308](#)]
7. Brown MT, Bussell JK. Medication adherence: WHO cares? *Mayo Clin Proc* 2011 Apr;86(4):304-314 [[FREE Full text](#)] [doi: [10.4065/mcp.2010.0575](#)] [Medline: [21389250](#)]
8. Cutler RL, Fernandez-Llimos F, Frommer M, Benrimoj C, Garcia-Cardenas V. Economic impact of medication non-adherence by disease groups: a systematic review. *BMJ Open* 2018 Jan 21;8(1):e016982 [[FREE Full text](#)] [doi: [10.1136/bmjopen-2017-016982](#)] [Medline: [29358417](#)]
9. Viswanathan M, Golin CE, Jones CD, Ashok M, Blalock SJ, Wines RC, et al. Interventions to improve adherence to self-administered medications for chronic diseases in the United States: a systematic review. *Ann Intern Med* 2012 Dec 4;157(11):785-795. [doi: [10.7326/0003-4819-157-11-201212040-00538](#)] [Medline: [22964778](#)]
10. Iuga AO, McGuire MJ. Adherence and health care costs. *Risk Manag Healthc Policy* 2014;7:35-44 [[FREE Full text](#)] [doi: [10.2147/RMHP.S19801](#)] [Medline: [24591853](#)]
11. Williams A, Kenyon K. eHealth Initiative. 2017. Issue brief: Medication adherence and health IT. URL: [https://www.ehidc.org/sites/default/files/resources/files/medicationadherence\\_and\\_hit\\_issue\\_brief\\_ONC.pdf](https://www.ehidc.org/sites/default/files/resources/files/medicationadherence_and_hit_issue_brief_ONC.pdf) [accessed 2019-06-13]
12. Carr-Lopez SM, Shek A, Lastimosa J, Patel R, Woelfel J, Galal S, et al. Medication adherence behaviors of Medicare beneficiaries. *Patient Prefer Adherence* 2014;8:1277-1284 [[FREE Full text](#)] [doi: [10.2147/PPA.S64825](#)] [Medline: [25258521](#)]



13. Centers for Medicare & Medicaid Services. 2018 Jul 12. CMS Proposes Historic Changes to Modernize Medicare and Restore the Doctor-Patient Relationship. URL: <https://www.cms.gov/newsroom/press-releases/cms-proposes-historic-changes-modernize-medicare-and-restore-doctor-patient-relationship> [accessed 2019-06-13]
14. Safavi K, Webb K, Kalis B. Accenture. 2019. Accenture 2019 Digital Health Consumer Survey. URL: <https://www.accenture.com/acnmedia/PDF-94/Accenture-2019-Digital-Health-Consumer-Survey.pdf> [accessed 2019-06-13]
15. Betts D, Korend L. Deloitte. 2018. Inside the Patient Journey: Three Key Touch Points for Consumer Engagement Strategies. URL: <https://www2.deloitte.com/insights/us/en/industry/health-care/patient-engagement-health-care-consumer-survey.html> [accessed 2019-06-13]
16. MDedge. 2019. VEText 1 Year Later - Still Growing. URL: <https://www.mdedge.com/fedprac/article/196293/health-policy/vetext-1-year-later-still-growing> [accessed 2019-10-07]
17. Veterans Affairs. 2019. VEText FAQs. URL: [https://www.va.gov/HEALTH/VEText\\_FAQs.asp](https://www.va.gov/HEALTH/VEText_FAQs.asp) [accessed 2019-10-07]
18. Thakkar J, Kurup R, Laba T, Santo K, Thiagalingam A, Rodgers A, et al. Mobile telephone text messaging for medication adherence in chronic disease: a meta-analysis. *JAMA Intern Med* 2016 Mar;176(3):340-349. [doi: [10.1001/jamainternmed.2015.7667](https://doi.org/10.1001/jamainternmed.2015.7667)] [Medline: [26831740](https://pubmed.ncbi.nlm.nih.gov/26831740/)]
19. Vervloet M, Linn AJ, van Weert JC, de Bakker DH, Bouvy ML, van Dijk L. The effectiveness of interventions using electronic reminders to improve adherence to chronic medication: a systematic review of the literature. *J Am Med Inform Assoc* 2012;19(5):696-704 [FREE Full text] [doi: [10.1136/amiainjnl-2011-000748](https://doi.org/10.1136/amiainjnl-2011-000748)] [Medline: [22534082](https://pubmed.ncbi.nlm.nih.gov/22534082/)]
20. Schwebel FJ, Larimer ME. Using text message reminders in health care services: a narrative literature review. *Internet Interv* 2018 Sep;13:82-104 [FREE Full text] [doi: [10.1016/j.invent.2018.06.002](https://doi.org/10.1016/j.invent.2018.06.002)] [Medline: [30206523](https://pubmed.ncbi.nlm.nih.gov/30206523/)]
21. Prayaga RB, Jeong EW, Feger E, Noble HK, Kmiec M, Prayaga RS. Improving refill adherence in medicare patients with tailored and interactive mobile text messaging: pilot study. *JMIR Mhealth Uhealth* 2018 Jan 30;6(1):e30 [FREE Full text] [doi: [10.2196/mhealth.8930](https://doi.org/10.2196/mhealth.8930)] [Medline: [29382623](https://pubmed.ncbi.nlm.nih.gov/29382623/)]
22. Zullig LL, Deschodt M, Liska J, Bosworth HB, de Geest S. Moving from the trial to the real world: improving medication adherence using insights of implementation science. *Annu Rev Pharmacol Toxicol* 2019 Jan 6;59:423-445. [doi: [10.1146/annurev-pharmtox-010818-021348](https://doi.org/10.1146/annurev-pharmtox-010818-021348)] [Medline: [30125127](https://pubmed.ncbi.nlm.nih.gov/30125127/)]
23. Laranjo L, Dunn AG, Tong H, Kocaballi AK, Chen J, Bashir R, et al. Conversational agents in healthcare: a systematic review. *J Am Med Inform Assoc* 2018 Sep 1;25(9):1248-1258 [FREE Full text] [doi: [10.1093/jamia/ocy072](https://doi.org/10.1093/jamia/ocy072)] [Medline: [30010941](https://pubmed.ncbi.nlm.nih.gov/30010941/)]
24. Bibault J, Chaix B, Nectoux P, Pienkowsky A, Guillemasse A, Brouard B. Healthcare ex Machina: Are conversational agents ready for prime time in oncology? *Clin Transl Radiat Oncol* 2019 May;16:55-59 [FREE Full text] [doi: [10.1016/j.ctro.2019.04.002](https://doi.org/10.1016/j.ctro.2019.04.002)] [Medline: [31008379](https://pubmed.ncbi.nlm.nih.gov/31008379/)]
25. Stein N, Brooks K. A fully automated conversational artificial intelligence for weight loss: longitudinal observational study among overweight and obese adults. *JMIR Diabetes* 2017 Nov 1;2(2):e28 [FREE Full text] [doi: [10.2196/diabetes.8590](https://doi.org/10.2196/diabetes.8590)] [Medline: [30291087](https://pubmed.ncbi.nlm.nih.gov/30291087/)]
26. Commission on Social Determinants of Health. World Health Organization. 2008. Closing the Gap in a Generation: Health Equity Through Action on the Social Determinants of Health. URL: [https://www.who.int/social\\_determinants/final\\_report/csdh\\_finalreport\\_2008.pdf](https://www.who.int/social_determinants/final_report/csdh_finalreport_2008.pdf) [accessed 2019-07-09]
27. Office of Disease Prevention and Health Promotion. Healthy People 2020. Social Determinants of Health. URL: <http://www.healthypeople.gov/2020/topics-objectives/topic/social-determinants-of-health> [accessed 2019-07-05]
28. Artiga S, Hinton E. Kaiser Family Foundation. 2018 May 10. Beyond Health Care: The Role of Social Determinants in Promoting Health and Health Equity. URL: <http://www.kff.org/disparities-policy/issue-brief/beyond-health-care-the-role-of-social-determinants-in-promoting-health-and-health-equity/> [accessed 2019-07-05]
29. Singh GK, Daus GP, Allender M, Ramey C, Martin E, Perry C, et al. Social determinants of health in the United States: addressing major health inequality trends for the nation, 1935-2016. *Int J MCH AIDS* 2017;6(2):139-164 [FREE Full text] [doi: [10.21106/ijma.236](https://doi.org/10.21106/ijma.236)] [Medline: [29367890](https://pubmed.ncbi.nlm.nih.gov/29367890/)]
30. Patel MR, Piette JD, Resnicow K, Kowalski-Dobson T, Heisler M. Social determinants of health, cost-related nonadherence, and cost-reducing behaviors among adults with diabetes: findings from the national health interview survey. *Med Care* 2016 Aug;54(8):796-803 [FREE Full text] [doi: [10.1097/MLR.0000000000000565](https://doi.org/10.1097/MLR.0000000000000565)] [Medline: [27219636](https://pubmed.ncbi.nlm.nih.gov/27219636/)]
31. Piette JD, Heisler M, Wagner TH. Cost-related medication underuse among chronically ill adults: the treatments people forgo, how often, and who is at risk. *Am J Public Health* 2004 Oct;94(10):1782-1787. [doi: [10.2105/ajph.94.10.1782](https://doi.org/10.2105/ajph.94.10.1782)] [Medline: [15451750](https://pubmed.ncbi.nlm.nih.gov/15451750/)]
32. National Academies of Sciences, Engineering, and Medicine. A Framework to Address the Social Determinants of Health. Washington, DC: The National Academies Press; 2016.
33. Reidel K, Tamblyn R, Patel V, Huang A. Pilot study of an interactive voice response system to improve medication refill compliance. *BMC Med Inform Decis Mak* 2008 Oct 9;8:46 [FREE Full text] [doi: [10.1186/1472-6947-8-46](https://doi.org/10.1186/1472-6947-8-46)] [Medline: [18845004](https://pubmed.ncbi.nlm.nih.gov/18845004/)]
34. Grice HP. Logic and conversation. In: Cole P, Morgan JL, editors. *Speech Acts*. New York: Academic Press; 1975:41-48.

## Abbreviations

**AI:** artificial intelligence  
**CAI:** conversational artificial intelligence  
**CMS:** Centers for Medicare and Medicaid Services  
**DOB:** date of birth  
**MLP:** multilayer perceptron  
**NLU:** natural language understanding  
**SDOH:** social determinants of health

*Edited by G Eysenbach; submitted 05.08.19; peer-reviewed by B Smith, K Davison, J Cornelius, S Brabyn, S Badawy, L McGoron; comments to author 27.08.19; revised version received 05.09.19; accepted 22.10.19; published 10.11.19*

*Please cite as:*

*Brar Prayaga R, Agrawal R, Nguyen B, Jeong EW, Noble HK, Paster A, Prayaga RS*

*Impact of Social Determinants of Health and Demographics on Refill Requests by Medicare Patients Using a Conversational Artificial Intelligence Text Messaging Solution: Cross-Sectional Study*

*JMIR Mhealth Uhealth 2019;7(11):e15771*

*URL: <http://mhealth.jmir.org/2019/11/e15771/>*

*doi: [10.2196/15771](https://doi.org/10.2196/15771)*

*PMID:*

©Rena Brar Prayaga, Ridhika Agrawal, Benjamin Nguyen, Erwin W Jeong, Harmony K Noble, Andrew Paster, Ram S Prayaga. Originally published in JMIR Mhealth and Uhealth (<http://mhealth.jmir.org>), 16.11.2019. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR mhealth and uhealth, is properly cited. The complete bibliographic information, a link to the original publication on <http://mhealth.jmir.org/>, as well as this copyright and license information must be included.